

University of Mississippi

eGrove

Annual Poster Session 2021

Annual Poster Session

10-19-2021

Liquid Chromatography- Tandem Mass Spectrometry (LC-MS-MS) as a detection method for Cocaine as a Performance Enhancing Drug

Isha Kalaria
University of Mississippi

Shrishti Dubey
University of Mississippi, sdubey@go.olemiss.edu

Follow this and additional works at: https://egrove.olemiss.edu/pharm_annual_posters_2021

 Part of the [Pharmacy and Pharmaceutical Sciences Commons](#)

Recommended Citation

Kalaria, Isha and Dubey, Shrishti, "Liquid Chromatography- Tandem Mass Spectrometry (LC-MS-MS) as a detection method for Cocaine as a Performance Enhancing Drug" (2021). *Annual Poster Session 2021*. 12. https://egrove.olemiss.edu/pharm_annual_posters_2021/12


This Book is brought to you for free and open access by the Annual Poster Session at eGrove. It has been accepted for inclusion in Annual Poster Session 2021 by an authorized administrator of eGrove. For more information, please contact egrove@olemiss.edu.

Liquid Chromatography- Tandem Mass Spectrometry (LC-MS-MS) as a detection method for Cocaine as a Performance Enhancing Drug

Made by- Shrishti Dubey, Isha Kalaria
Department of Pharmaceutics and Drug Delivery, School of Pharmacy
The University of Mississippi

INTRODUCTION

In competitive sports, doping is the use of banned athletic performance-enhancing drugs by athletic competitors. Among the most popular PEDs are anabolic steroids, human growth hormone, erythropoietin (EPO), beta-blockers, stimulants and diuretics.



Sports with the greatest numbers of anti-doping rules violations in 2015.

World Anti-Doping Authority (WADA) : It works to coordinate the fight against doping in sport internationally WADA prevents the use of Performance enhancing drugs because: 1) it is or potentially is harmful to the athlete's health, and 2) it violates the spirit of sport.

WADA has published an annual List of Prohibited Substances and Methods

Analytical techniques look at not only banned substances but metabolites

Samples that can be tested include blood, saliva and urine, with a wide range of techniques being used: GC-MS, LC-MS, LC-MS-MS, Biomarker tests, Isoform Differential Immunoassay Tests, RNA Testing, GC/C/IRMS, Flow Cytometry, Isoelectric Focusing.

METHOD

Liquid Chromatography - Tandem Mass Spectrometry (LC-MS-MS)

It is a powerful analytical technique that combines the separating power of liquid chromatography with the highly sensitive and selective mass analysis capability of triple quadrupole mass spectrometry.

ADVANTAGES


- ✓ Quicker and less extensive extraction procedures
- ✓ Simultaneous multianalyte method.
- ✓ Highly selective, specific and gives accurate results within a short span of time
- ✓ Better sensitivity than GC.
- ✓ No derivatization needed.

DISADVANTAGES

- ✓ Expensive.
- ✓ Not portable.
- ✓ Experienced Technicians required.

INSTRUMENTATION

- MS-MS is the combination of two mass analyzer in one mass spectra instrument.
- The first mass filters for precursor ion followed by a fragmentation of the precursor ion with high energy and example is nitrogen gas.
- A second mass analyzer is then filtering the product ions, generated by the fragmentation. This is usually done in a triple quadrupole MS(QQQ) or a(QTOF).



MASS SPECTROMETRY OF COCAINE




Figure 1. Product ion spectra of protonated COC (A), BE (B), EME (C), and ECG (D) at 21 eV collision energy.

COCAINE

Benzoyllecgonine (BZE), a demethylation product of cocaine (COC) and the major metabolite found in humans, does not exhibit any pharmacological activity. It has wide toxicological interest however, due to its long half-life, and remains detectable in biological fluids considerably longer than COC.

Major Metabolites of Cocaine:

- Benzoylecgonine (BE)
- Ecgonine methyl ester (EME)
- Ecgonidine

CONCLUSION

LC- MS-MS is one of the best techniques to find out the concentration of cocaine from samples. The peaks obtained from the Mass Spectrometer helps in determining that the compound that we are detecting is present in the sample.

REFERENCES

1. Sveshnikova, N., Yuan, T., Warren, J.M. et al. Development and validation of a reliable LC-MS/MS method for quantitative analysis of usnic acid in *Cladonia uncialis*. BMC Res Notes 12, 550 (2019). <https://doi.org/10.1186/s13104-019-4580-x>
2. Perez ER, Knapp JA, Horn CK, Stillman SL, Evans JE, Arfsten DP. Comparison of LC-MS-MS and GC-MS Analysis of Benzodiazepine Compounds Included in the Drug Demand Reduction Urinalysis Program. J Anal Toxicol. 2016;40(3):201-207. doi:10.1093/jat/bkv140
3. Analysis of cocaine, benzoylecgonine, ecgonine methyl ester, and ecgonine by high-pressure liquid chromatography-API mass spectrometry and application to a short-term degradation study of cocaine in plasma
4. Klingmann, A., Skopp, G. and Aderjan, R., 2021. Analysis of Cocaine, Benzoylecgonine Ecgonine Methyl Ester, and Ecgonine by High-Pressure Liquid Chromatography-API Mass Spectrometry and Application to a Short-Term Degradation Study of Cocaine in Plasma.
5. En.wikipedia.org. 2021. Liquid chromatography-mass spectrometry - Wikipedia. [online] Available at: <https://en.wikipedia.org/wiki/Liquid_chromatography%E2%80%93mass_spectrometry> [Accessed 8 October 2021].
6. Shpak, A., Appolonova, S. and Semenov, V., 2021. Validation of Liquid Chromatography-Electrospray Ionization Ion Trap Mass Spectrometry Method for the Determination of Mesocarb in Human Plasma and Urine.
7. Shpak, A., Appolonova, S. and Semenov, V., 2021. Validation of Liquid Chromatography-Electrospray Ionization Ion Trap Mass Spectrometry Method for the Determination of Mesocarb in Human Plasma and Urine.